


National
Library
of Medicine

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Search

PubMed

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

Display

Abstract

Show

20

Sort by

Send to

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: Biophys J. 2004 Dec;87(6):3842-9. Epub 2004 Sep 17. Related Articles, LinksFull text article at
www.biophysj.org**Protein translocation through anthrax toxin channels formed in planar lipid bilayers.****Zhang S, Udho E, Wu Z, Collier RJ, Finkelstein A.**

Department of Microbiology and Molecular Genetics, Harvard Medical School, Boston, Massachusetts, USA.

The 63-kDa fragment of the protective antigen (PA) component of anthrax toxin forms a heptameric channel, (PA63)₇, in acidic endosomal membranes that leads to the translocation of edema factor (EF) and lethal factor (LF) to the cytosol. It also forms a channel in planar phospholipid bilayer membranes. What role does this channel play in the translocation of EF and LF? We report that after the 263-residue N-terminal piece of LF (LFN) binds to its receptor on the (PA63)₇ channel and its N-terminal end enters the channel at small positive voltages to block it, LFN is translocated through the channel to the opposite side at large positive voltages, thereby unblocking it. Thus, all of the translocation machinery is contained in the (PA63)₇ channel, and translocation does not require any cellular proteins. The kinetics of this translocation are S-shaped, voltage-dependent, and occur on a timescale of seconds. We suggest that the translocation process might be explained simply by electrophoresis of unfolded LFN through the channel, but the refolding of the N-terminal half of LFN as it emerges from the channel may also provide energy for moving the rest of the molecule through the channel.

PMID: 15377524 [PubMed - indexed for MEDLINE]

Display

Abstract

Show

20

Sort by

Send to

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Jun 27 2005 04:57:20